


Figure 1. Treatment periods and insecticide choices for management of San Jose scale. Rates are listed as amount of material/100 gallon.



① a	② b	③ c	④ d
<b>GREEN – TIP</b>	<b>HALF-INCH-GREEN</b>	<b>1<sup>st</sup> SUMMER BROOD (crawlers)</b>	<b>2<sup>nd</sup> SUMMER BROOD (crawlers)</b>
		<i>Apply @ 500 DD from 1 March + same 14d later</i>	<i>Apply @ 1451 DD from 1 March + same 14d later</i>
3% oil	Use either: 2% oil Lorsban 4E (16 oz) Lorsban + oil Supracide 2E (32 oz) Supracide 2E + oil	Use either: Esteem (1 oz) + oil (2%) Assail (1 oz) + oil (2%) Provado 1.6F (2 oz) + oil (2%)	Use either: Esteem (1 oz) + oil (2%) Assail (1 oz) + oil (2%) Provado 1.6F (2 oz) + oil (2%)

a. In low risk orchards, this treatment alone may be sufficient.

b. Treatment during this period is recommended if harvest damage was noted the previous season.

c. Treatment during this period may be necessary if bark infestations and/or fruit damage was noted the previous season.

d. Examine fruit (mid-July) to determine need for late-season treatments.