

# SCAFFOLDS Fruit Journal, Geneva, NY

Volume 22, No. 20

Update on Pest Management and Crop Development

August 5, 2013

---

## COMING EVENTS

	43°F	50°F
Current DD accumulations		
(Geneva 1/1-8/5):	2376	1641
(Geneva 1/1-8/5/2012):	2855	2006
(Geneva "Normal" for this date):	2400	1617
(Geneva 1/1-8/12 predicted):	2570	1786
(Highland 1/1-8/5/2013):	2752	1937

## Upcoming Pest Events – Ranges (Normal +/- Std Dev):

### American plum borer

2nd flight peak ..... 2002-2856 1347-1785

Apple maggot flight peak..... 2103-2657 1408-1838

Codling moth 2nd flight peak..... 1931-2735 1278-1892

### Comstock mealybug

2nd gen. crawlers emerge..... 2234-2624 1505-1781

### Comstock mealybug

2nd gen. crawlers peak ..... 2380-2624 1658-1737

### Lesser appleworm

2nd flight peak ..... 2131-3105 1422-2156

### Obliquebanded leafroller

2nd flight peak .....	2593-3011	1758-2098
Oriental fruit moth		
2nd flight subsides .....	2069-2567	1376-1794
Redbanded leafroller		
2nd flight subsides .....	2182-2742	1471-1891
San Jose scale 2nd flight peak....	2128-2500	1434-1750
Spotted tentiform leafminer		
3rd flight begins .....	2257-2655	1512-1844

TRAP CATCHES (Number/trap/day)

Geneva

	7/25	7/29	8/2	8/5
Redbanded Leafroller	0.3	0.0	0.0	0.0
Spotted Tentiform Leafminer	9.0	4.1	6.0	2.5
Oriental Fruit Moth	0.0	0.0	0.3*	0.5
San Jose Scale	44.8	57.5	198	173
Codling Moth	0.0	0.0	1.3*	1.5
American Plum Borer	0.2	0.5	0.5	0.3
Lesser Peachtree Borer	0.0	0.1	0.0	0.0
Obliquebanded Leafroller	0.0	0.0	0.0	0.1*
Dogwood Borer	1.2	0.8	-	0.9
Apple Maggot	0.3	0.1	0.1	1.0

Highland (Peter Jentsch)

7/15	7/22	7/29	8/5
------	------	------	-----

Redbanded Leafroller	2.1	0.6	0.2	0.4
Spotted Tentiform Leafminer	10.5	28.8	12.1	24.0
Oriental Fruit Moth	1.2	1.3	0.0	0.7
Lesser Appleworm	0.2	0.9	0.4	0.4
Codling Moth	0.6	0.3	1.3	1.3
Obliquebanded Leafroller	0.2	0.2	0.2	0.2
San Jose Scale	44.5	148	16.6	1.0
Apple maggot	0.1	0.5	0.2	0.1

\* = 1st catch

## Pest Focus

Geneva: Obliquebanded Leafroller 2nd flight began today, 8/5.

Oriental Fruit Moth 3rd flight and Codling Moth 2nd flight began 8/2.

Highland: San Jose Scale crawler emergence predicted for 8/19 (using 600-700 DD50 from biofix.)

## ORCHARD RADAR DIGEST

**[Box Text: ECHOES]**

## **Geneva Predictions:**

### Codling Moth

Codling moth development as of August 4: 2nd generation adult emergence at 63% and 2nd generation egg hatch at 24%.

2nd generation 30% CM egg hatch: August 7 (= target date where one spray needed to control 2nd generation CM).

### White Apple Leafhopper

2nd generation WAL found on apple foliage: August 6.

## **[Section: INSECTS]**

### SWD UPDATE

(Julie Carroll, NYS IPM Program, Geneva;  
jec3@cornell.edu)

## **[Box text: GETTING SPOTTY]**

A single female SWD was caught on July 29 in one of two traps in a sweet cherry block in Ontario Co. that had been harvested already.

SWD were caught in traps collected on July 29 that were set in and near high tunnel raspberries in Tioga County.

A single male SWD was caught on July 30 in one of two traps set in a raspberry planting in Niagara County.

For your reference, a quick guide to the insecticides labeled and available for use against SWD in the following crops have been posted on the Cornell Fruit website:

### **Stone Fruits and Grapes**

<http://www.fruit.cornell.edu/spottedwing/pdfs/TreeFruitGrapeSWDinsecticides2013.pdf>

### **Berry Crops**

<http://www.fruit.cornell.edu/spottedwing/pdfs/UpdatedLabeledInsecticidesNY-SWD-Final.pdf>

## **HUDSON VALLEY PEST UPDATE**

(Peter Jentsch, Entomology, HighlAnd)

**[Box text: DOWN IN THE VALLEY]**

### **Apple Maggot**

Our first apple maggot adult fly was captured on the 8th of July this season, somewhat later than what we've observed in Highland in previous years. Apple maggot adult flight numbers have been relatively low in traps over the past few weeks, with threshold reached for a first application on the early varieties such as Ginger

Gold on July 2. Given moist soil conditions, it's likely higher numbers of AM will emerge over the next few weeks. (Click for AM options

<http://ipmguidelines.org/TreeFruits/Chapters/CH11/default-5-8.aspx>).

In years of heavy rainfall late in the season, questions arise regarding the efficacy of

residual insecticide for AM control, related to rain

events. John Wise, Michigan State University,

Department of Entomology, wrote a very nice piece on

the "Rainfast characteristics of fruit crop insecticides"

that might help to answer questions regarding this

topic.

([http://msue.anr.msu.edu/news/rainfast\\_characteristics\\_of\\_fruit\\_crop\\_insecticides](http://msue.anr.msu.edu/news/rainfast_characteristics_of_fruit_crop_insecticides))

## **Brown Marmorated Stink Bug (BMSB) Update**

We are seeing large numbers of 5th instar nymphs in traps placed along orchard perimeters in Orange,

Ulster, Dutchess and Columbia Counties. The 1st

generation adults are now beginning to move into tree

fruit in a few sites, in low numbers. Peaches and early

apples are very susceptible at this time. Highest

numbers of nymphs have been observed in Marlboro.

A low percentage of fruit injury has been observed

along the perimeters of apple blocks in a Columbia

County orchard this past week.

Strategies for BMSB management should begin with perimeter scouting along wooded edges and hedgerows of orchards to optimize your efforts. Knowing the various BMSB life forms - eggs, nymphs and adults - is important during decision making, as there are many different types of stink bug, with similar physical characteristics, present this time of year. Applications should only be made after confirmed BMSB sightings in the orchard have been made. (Click for BMSB identification

<http://hudsonvf.cce.cornell.edu/scouting%20reports/BMSB%20Project/BMSB%20Update%20in%20New%20York%20State%20%20Fall%202011.pdf> .

Applications should be made in orchard blocks once the first BMSB has been found on fruit trees. During the 2012 growing season, the adults were seen moving into apple in late August with significant injury observed by mid-September. Given the elusive nature of this pest, low populations observed during scouting can equate to relatively high in-orchard numbers that can cause severe fruit injury. Since this insect causes the greatest damage along the perimeter of apple and peach orchards, applications directed along the first 90 feet of outer orchard rows will reduce the majority of

the damage. Control options to manage BMSB can be found here

(<http://hudsonvf.cce.cornell.edu/scouting%20reports/BMSB%20Project/NY%20BMSB%20Insecticides%20&%20Efficacy%207-2013.pdf>).

## **[Section: GENERAL INFO]**

### EVENT ANNOUNCEMENTS

#### **[Box text: IT'S A DATE]**

#### WAYNE COUNTY FRUITGROWER TOUR

Wednesday, August 7, from 10:00 am

Registration and 1st stop at Empire Fruit Packers & Lake Country Storage, 10529 Ridge Rd., Wolcott, NY

Sponsored by agr.assistance, this large, informative and entertaining tour is in its 15th year, and will feature presentations on apple storage; Honeycrisp planting strategies; solar protection; powdery mildew control; weed control; frost protection; deer and bird control; managing biennial apple varieties; fire blight control; internal worm control programs, plus much more.

Door prizes, lunch, some levity, a BBQ/clambake dinner with a live band, growers and industry representatives



from NY and surrounding states — tough to beat on a midsummer day. Free attendance.

Contact Lindsay LaMora (585-734-8904; [lindsaylamora@agrassistance.com](mailto:lindsaylamora@agrassistance.com)) for RSVP pre-registration and tour information.

## CORNELL FRUIT PEST CONTROL FIELD DAYS

The N.Y. Fruit Pest Control Field Days will take place during Labor Day week on Sept. 4 and 5 this year, with the Geneva portion taking place first (Wednesday Sept. 4), and the Hudson Valley installment on the second day (Thursday Sept. 5). Activities will commence in Geneva on the 4th, with registration, coffee, etc., in the lobby of Barton Lab at 8:30 am. The tour will proceed to the orchards to view plots and preliminary data from field trials involving new fungicides, bactericides, miticides, and insecticides on tree fruits and grapes. It is anticipated that the tour of field plots will be completed by noon. On the 5th, participants will register at the Hudson Valley Laboratory starting at 8:30, after which they will view and discuss results from field trials on apples and other fruit crops. Although Dave Rosenberger did not run sponsored trials this year, he will be reporting on four field trials involving efficacy of copper products in green tip sprays, potential of Blossom Protect to russet fruit when

applied to control blossom blight, scab and rust control with ProPhyt and AgriFos, and efficacy of summer fungicides applied after sooty blotch and flyspeck have become established. No pre-registration is required for either event.

---

This material is based upon work supported by Smith Lever funds from the Cooperative State Research, Education, and Extension Service, U.S. Department of Agriculture. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.

Scaffolds is published weekly from March to September by Cornell University -- NYS Agricultural Experiment Station (Geneva), and Ithaca -- with the assistance of Cornell Cooperative Extension. New York field reports welcomed. Send submissions by 2 p.m. Monday to:

Scaffolds Fruit Journal

Editors: A. Agnello, D. Kain

Dept. of Entomology, NYSAES

630 W. North St.

Geneva, NY 14456-1371

Phone: 315-787-2341 FAX: 315-787-2326

E-mail: ama4@cornell.edu

Online at

<<http://www.scaffolds.entomology.cornell.edu/index.html>>