

SCAFFOLDS Fruit Journal, Geneva, NY

Volume 25, No. 23

Update on Pest Management and Crop Development

September 6, 2016

COMING EVENTS

	43°F	50°F
Current DD* accumulations	3354	2360
(Geneva 1/1-9/6):	3082	2136
(Geneva 1/1-9/6/2015):	3082	2136
(Geneva "Normal"):	3235	2237
(Geneva 1/1-9/12, predicted):	3531	2495
(Highland 1/1-9/6):	3918	2794

Upcoming Pest Events – Ranges (Normal +/- Std Dev):

American plum borer

2nd flight subsides2927-3353 2018-2372

Codling moth

2nd flight subsides2846-3462 1923-2447

Lesser appleworm

2nd flight subsides2794-3488 1918-2422

Lesser peachtree borer

flight subsides2996-3446 2017-2433

Obliquebanded leafroller

2nd flight subsides3108-3468 2126-2448

Oriental fruit moth

3rd flight subsidies	2928-3412	1978-2310
Redbanded leafroller		
3rd flight subsidies	3124-3436	2142-2422
Spotted tentiform leafminer		
3rd flight subsidies	3244-3480	2258-2462
*[all DDs Baskerville-Emin, B.E.]		

TRAP CATCHES

Geneva (Number/trap)

	8/25	8/29	9/2	9/6
Redbanded Leafroller	19.0	6.5	4.0	1.0
Spotted Tentiform Leafminer	24.5	5.5	6.0	7.0
Oriental Fruit Moth	4.0	2.0	5.5	5.5
Lesser Apple Worm	2.0	0.0	1.5	0.5
Codling Moth	7.5	2.0	1.5	2.0
American Plum Borer	0.0	0.0	0.0	0.0
Lesser Peachtree Borer	1.5	3.0	1.0	0.5
Obliquebanded Leafroller	0.0	0.5	0.5	0.0
Pandemis Leafroller	0.5	0.0	0.5	0.5
Dogwood Borer	0.0	0.0	0.5	0.0
Peachtree Borer	3.0	0.0	0.0	0.0
Apple Maggot	0.3	1.7	2.0	1.0

Highland (Peter Jentsch)

	8/15	8/22	8/29	9/6
Redbanded Leafroller	38.5	18.0	36.5	33.5

Spotted Tentiform Leafminer	74.0	13.0	13.5	38.5
Oriental Fruit Moth	14.0	8.5	11.0	18.5
Lesser Appleworm	26.0	19.5	36.5	35.5
San Jose Scale	9.0	481.5	107.5	73.0
Codling Moth	41.5	20.0	12.0	13.5
Obliquebanded Leafroller	14.5	11.5	11.0	10.0
Dogwood Borer	8.5	4.5	1.5	1.0
Brown Marmorated Stink Bug	0.0	0.0	0.0	0.0
Apple Maggot	14.8	4.8	8.0	7.3

[Section: DISEASES]

APPLE TRUNK PROBLEMS CAN REDUCE PROFITABILITY
(Dave Rosenberger, Plant Pathology, Highland;
dar22@cornell.edu)

[Box text: FROM THE GROUND UP]

Over the past 15 years, I have become increasingly aware of the many factors that can adversely impact the growth and health of apple tree trunks. I've also vacillated between amusement and despair on various farm tours where horticulturists have expounded at length on how to prune, train, and improve fruit quality on young trees in blocks where a more "down toward earth" view would have revealed trunk problems that

were almost certain to limit profitability of the block regardless of how trees were pruned or trained.

Unfortunately, trunk problems often evolve from a variety of interacting stresses, pathogens, and/or insects. The initial causes are sometimes difficult to pinpoint, and therefore solutions to these problems remain elusive. Following is a list of problems that I have observed along with my subjective perspectives on potential contributing factors and possible solutions. I'm certain that other folks will have additional and divergent opinions about causes and solutions for many of these problems, but I hope that my mentioning them will at least stimulate useful thinking and discussion.

1. Rootstock blight killing young trees: Fire blight that gets into M.9 or M.26 rootstocks in trees less than 8 or 10 years old will almost always be lethal (Fig. 1). The key to avoiding tree loss to this disease is to order all new trees on rootstocks other than the highly susceptible M.9 or M.26. If trees are propagated and planted on either of these highly susceptible rootstocks, then growers should expect to spend extra time and money protecting blossoms with antibiotics and/or copper sprays during at least the first eight years after trees are planted, and probably for the lifetime of the

orchard. Given our changing climate and the fact that many new cultivars are very blight susceptible, one must assume that blight epidemics can occur anywhere in North America, even if some regions rarely had fire blight in the past.

2. Excessive production of root suckers in young trees:

Root suckers (Fig. 2) are bad because they are difficult to manage, may prevent full coverage of trunk sprays applied to control borers, and can absorb and translocate herbicides such as glyphosate. If the rootstock is M.9 or M.26, actively growing suckers provide an entry point for fire blight that will quickly kill those rootstocks. Perhaps it is only my idealized memories of the past, but my impression is that root suckers are more prevalent in young trees and are developing in larger numbers than was the case 30 years ago. I don't know if changing nursery practices or changing cultural practices in the orchard are contributing factors. (Some folks have suggested that every time a sucker is burned down with glyphosate, all of its cousins emerge to seek revenge.) Thirty years ago, the suggested planting depth for new trees required graft unions to be two inches above the soil line after trees had settled. Because of problems with scion rooting in some orchards where graft unions

ended up at or below ground level, the suggested planting height gradually changed from 2 inches to 4 inches to 6 inches, and trees that fail to settle as expected after planting may end up with 8 or 10 inches of rootstock exposed. Increasing the length of the rootstock above ground generally reduces vigor of the scion and may promote production of more root suckers. We know that some rootstocks produce more suckers than others. I don't have a solution for avoiding or managing suckers, but everyone should recognize that suckers pose a threat to tree health.

3. Burr knots: Burr knots have always posed problems on some rootstocks, but they become even more problematic on trees that are planted with 6 to 8 inches of the rootstocks exposed. Trees with two or three strategically located burr knots often have distorted trunks (Fig. 3). Burr knots are especially attractive to dogwood borers. Perhaps there are nursery practices that can reduce the incidence of burr knots, but growers can help to minimize this problem by avoiding graft unions that are higher than necessary. In some cases, especially if the planting of grass in row middles is delayed until the fall after trees are planted, trees with excessively high graft unions could be bermed slightly after trees have settled, thereby reducing risks

that burr knots will develop on the excessively exposed rootstocks.

4. Herbicide injury on trunks: Initially I believed that glyphosate was the cause of most trunk damage on apple trees (see <http://www.scaffolds.entomology.cornell.edu/2004/040615.html> from 14 June 2004). Glyphosate uptake through trunks is still contributing to trunk injury, but it is becoming apparent that other herbicides, while not translocated throughout the tree as is the case for glyphosate, may still be causing localized damage to trunks (Fig. 4). Gramoxone, glufosinate (Rely), and perhaps other products can be nearly as injurious as glyphosate, especially if trees receive multiple applications each year. I suspect that many herbicides cause a slight desiccation of young bark that is hit by the sprays. Repeated desiccation over several years reduces growth of the lower trunk and encourages infection by *Botryosphaeria dothidea*, the white rot pathogen, as explained in the next paragraph. Repeated desiccation sometimes results in decreased trunk diameters just above the graft unions (Fig. 5). Painting trunks with a white latex paint (Fig. 6) appears to protect the trunks to some degree (research needed!). However, it is very difficult to get good paint

coverage on the exposed portions of the rootstocks that are often roughened by burr knots and flaky bark. In some cases, we are now seeing rootstocks dying from apparent herbicide damage where the trunks that were protected by white paint appear healthy.

Cultivars and rootstocks seem to differ significantly in their susceptibility to damage from herbicides. The only solution here seems to be to paint trunks on young trees, keep graft unions as low as is practical without risking scion rooting, and minimize exposure of trunks to ANY and ALL herbicides, especially on trees less than 6 or 8 years old. Trunk guards (such as milk carton sleeves) placed around young trees can minimize herbicide contact with the trunks, but if trunk guards are left in place throughout the year, they may eventually serve to attract and protect borers.

5. Drought injury on trunks: Apple trees subjected to drought stress during summer will eventually be attacked by *B. dothidea*, an opportunistic fungus that is resident in older bark and that can damage (Fig. 7) and even kill the bark and cambium on trees that are water stressed. When periods of water stress are limited, *B. dothidea* may kill only the outer bark, which then peels off in large flakes to reveal healthy, relatively green bark beneath the flakes. Recently, Dr. Srdjan Acimovic,

plant pathologist at the Hudson Valley Lab, suggested that flaking bark, which is more prevalent on some rootstocks than on others, might expose thin green bark beneath the flakes to herbicides, thereby allowing injury of the green bark that would not occur in the absence of the flaky bark. Thus, herbicide desiccation of bark may favor infection and invasion of *B. dothidea* that then causes more flaky bark that exposes underlying green bark to herbicide damage as the flakes are shed, thus creating a cycle that allows continued damage in older trees. The solution for avoiding drought stress is to ensure that new plantings can be irrigated. I am increasingly convinced that trees on M.9 rootstocks should not be planted on any ground that cannot be irrigated because this rootstock will not survive the drought stresses that will likely become more prevalent throughout the northeastern United States as a result of climate change over the next 20 or 30 years. For those without access to (enough) irrigation water, the million-dollar question is, which of the new rootstocks is best suited for surviving drought stress, and at what spacing should they be used on non-irrigated sites? I'm not sure anyone knows, but I'm fairly certain G.11 will be no better than M.9 on droughty sites because I have already seen severe *B.*

dothidea on drought-stressed trees on G.11 at the Hudson Valley Lab.

6. Low pH causing measles (Fig. 8): Most growers are doing a good job of adjusting soil pH prior to planting, but we still see occasional plantings where tree health is jeopardized by low pH. Bringing up soil pH after planting is much more difficult than if it is done prior to planting.

7. Latent viruses: Apple trees can carry four latent viruses that are transmitted only via propagation or by root grafting from adjacent trees. The latent viruses are stem pitting, stem grooving, apple chlorotic leaf spot, and apple mosaic. These latent viruses generally do not cause any visible symptoms or direct tree decline, although they can reduce yield in some cultivars. In the earlier part of the 20th century, before these viruses had been identified, apple growers sometimes used Virginia Crab as a rootstock. When apple stem pitting and/or apple stem grooving was introduced into the Virginia crab rootstock by grafting a scion cultivar that contained these viruses, the rootstock would develop severe pitting (Fig. 9) that stunted the trees and made them unproductive. Clean nursery stock programs initiated during the 1950s and 1960s pretty much

eliminated these virus problems, but those programs gradually were abandoned due to lack of interest and funding. Now tree decline from latent viruses is re-emerging as a commercial problem. It behooves growers to request and insist upon trees that are propagated using virus-free scion material, so as to avoid tree losses several years after planting.

8. Borer problems: Dogwood borer (an old problem) and black stem borer (a newer problem) can devastate trunks on young trees. Dogwood borer is favored by the presence of burr knots that provide entry points and by flaky bark and abundant root suckers, both of which can provide protection from both predators and insecticide sprays. Changing pesticide registrations may make management of these borers more difficult in the future. Failure to monitor carefully for borer problems and apply controls when borers first appear can rapidly lead to tree decline and other trunk problems. There is some question about whether or not borer damage provides entry points for contact herbicides that then contribute to further trunk injury.

Ultimately, good tree health is heavily dependent on having healthy tree trunks. Growers who are investing up to \$30,000 per acre in new plantings should be

asking tough questions about rootstocks, virus contamination of budwood, recommended planting depths, and why some lots of trees have so many burr knots or produce so many suckers within a few years after planting. Perhaps no one has all of the answers at this point, but we can't expect much progress if no one is thinking about how to improve the key structures (apple trunks) that support all apple production systems.

[Section: INSECTS]

TRAPPED OUT

(Art Agnello, Entomology, Geneva; ama4@cornell.edu)

[Box Text: COOLING IT]

With this issue, Scaffolds ceases publication for the season; we expect to start up again next March. In March, as usual, we'll send out an email to all current subscribers to verify addresses for next year's mailing list. Our thanks to all of you who have sent comments, suggestions, and articles our way, a practice we hope you'll continue. As a wrap-up, here's our annual summary of the year's pheromone trap results and an Index of Volume 25, 2016 of Scaffolds Fruit Journal.

KEY = GFW - Green Fruitworm; RBLR - Redbanded Leafroller; STLM - Spotted Tentiform Leafminer; OFM - Oriental Fruit Moth (in apples); LAW - Lesser Appleworm; CM - Codling Moth; APB - American Plum Borer (in peach); LPTB - Lesser Peachtree Borer (in peach); DWB - Dogwood Borer; PL - Pandemis Leafroller; OBLR - Obliquebanded Leafroller; PTB - Peachtree Borer; AM - Apple Maggot; * - first catch of the generation.

Geneva Pest Trapping Results - Avg/Trap

DATE	GFW	RBLR	STLM	OFM	LAW	CM	SJS	APB	LPTB	OBLR
3/25	1.0*									
3/28	8.5									
4/7	14.0									
4/11	16.0	0.0								
4/15	11.5	0.5*	0.0							
4/18	9.5	23.0	1.0*							
4/22	19.5	114.5	62.5							
4/25	18.0	26.5	2.5							
4/29	6.0	9.5	0.0	0.0						
5/2	1.5	14.5	12.0	3.5*						
5/5	0.5	37.0	64.0	5.5						
5/9	3.0	11.5	40.0	5.5						
5/12	0.0	31.0	22.0	31.0						
5/16	0.0	21.0	18.0	19.5						
5/20	0.5	19.5	20.5	11.0	0.0	0.5*				
5/23	0.0	26.5	36.0	31.5	0.5*	2.5				

5/26		21.5	15.0	29.5	0.0	11.5	3.0*	1.0*	12.5*			
5/31		10.0	3.0	29.5	4.5	15.5	3.0	0.0	13.0	3.0*		
DATE	RBLR	STLM	OFM	LAW	CM	SJS	LPTB	OBLR	DWB	PL	PTB	AM
6/3	0.0	1.0	5.0	2.5	5.0	0.5	18.5	1.5	2.5*	1.0*		
6/6	0.0	1.5	4.5	3.0	2.5	0.0	8.0	1.0	2.0	5.5		
6/9	0.0	0.0	0.0	0.0	0.5	0.5	0.5	1.0	0.5	2.0		
6/13	0.0	1.5	0.0	0.5	1.0	0.0	4.5	2.0	-	31.0		
6/16	0.0	9.0	0.0	2.0	2.0		3.0	3.0	1.5	1.5	1.0*	
6/21	12.5*	19.5	0.0	1.0	7.5		3.5	27.0	5.0	8.5	7.0	
6/24	1.5	31.5	0.5	1.0	0.0		0.0	8.5	8.0	1.5	4.5	
6/27	9.0	78.0	4.0	1.5	11.5		0.5	7.0	15.5	3.0	1.0	
6/30	8.0	123.0	0.0	0.0	2.0		1.5	2.0	30.0	2.0	2.5	
7/5	8.0	158.0	1.0	0.0	3.5		0.5	0.0	23.5	1.0	3.0	0.0
7/8	6.5	182.5	2.5	0.5	1.0		0.0	3.0	8.0	0.0	9.5	0.3*
7/11	2.5	42.5	3.5	0.0	0.0		0.0	0.5	8.0	0.0	3.0	0.0
7/15	19.5	73.0	0.0	0.0	0.5		0.5	0.0	0.0	0.0	6.0	0.0
7/18	9.5	53.0	0.0	0.0	4.5		0.0	0.0	0.5	0.0	0.0	0.0
7/25	11.0	31.0	0.0	1.0	24.0		0.0	0.0	0.0	0.0	0.5	0.7
7/28	2.5	26.5	0.0	1.0	5.0		1.0	0.0	1.0	0.0	0.5	0.3
8/1	0.0	112.0	4.5*	5.0	8.0		0.0	0.0	0.0	0.0	4.0	3.3
8/5	1.5	78.5	5.0	2.5	5.5		1.5	0.0	0.5	0.0	4.0	3.0
8/8	2.0	48.0	1.0	0.0	4.5		4.0	0.5	0.0	0.5	1.0	0.0
8/12	8.5	57.0	9.5	2.0	7.5		5.5	1.0	0.0	0.5	4.0	0.3
8/15	25.0	39.0	4.0	1.5	11.0		3.5	1.0	1.0	0.0	5.5	0.3
8/19	19.0	40.0	3.0	4.5	8.5		4.0	0.0	0.0	1.0	2.0	0.3
8/25	19.0	24.5	4.0	2.0	7.5		1.5	0.0	0.0	0.5	3.0	0.3
8/29	6.5	5.5	2.0	0.0	2.0		3.0	0.5	0.0	0.0	0.0	1.7
9/2	4.0	6.0	5.5	1.5	1.5		1.0	0.5	0.5	0.5	0.0	2.0
9/6	1.0	7.0	5.5	0.5	2.0		0.5	0.0	0.0	0.5	0.0	1.0

HUDSON VALLEY INSECT KEY = GFW - Green Fruitworm; RBLR - Redbanded Leafroller; STLM - Spotted Tentiform Leafminer; OFM - Oriental Fruit Moth (in apples); LAW - Lesser Appleworm; CM - Codling Moth; SJS - San Jose scale; DWB - Dogwood borer; OBLR - Obliquebanded Leafroller; AM - Apple Maggot; * - first catch of the generation.

Hudson Valley (Highland) Pest Trapping Results - Avg/Trap

DATE	GFW	RBLR	STLM	OFM	LAW	DWB	SJS	CM	OBLR	AM
3/14		1.0*								
3/21	0.4*	0.5								
3/28	7.0	26.0								
4/4	0.0	4.5	20.5*							
4/11	1.0	12.5	6.5							
4/18	0.5	45.0	136.5							
4/25	1.5	151.0	678.0							
5/2	0.5	28.0	89.0	13.0*						
5/9	1.0	18.0	53.0	19.0	2.0*					
5/16	0.0	19.0	25.5	27.5	17.5	1.0*	513.5*			
5/23		3.0	17.5	4.5	2.0	0.0	2.0	35.0*		
5/31		1.0	1.5	0.5	5.0	1.0	32.0	71.0	4.5*	
6/6		0.0	60.0	1.0	4.0	2.0	0.0	44.5	21.0	
6/13		0.5	96.0	0.0	1.0	1.1	0.0	35.5	41.5	
6/20		8.5	327.0	0.5	7.5	0.0	0.5	48.5	62.5	
6/27		39.0	255.5	3.5	21.5	1.5	10.0	31.5	63.0	
7/5		45.0	214.5	4.5	25.0	9.5	1.5	7.5	22.0	
7/11		21.5	119.5	10.0	10.0	5.0	2.5	32.5	11.0	2.8*
7/18		14.5	60.5	3.5	4.0	5.5	15,680	69.0	5.5	4.3

7/25	3.5	151.0	1.5	9.0	2.5	3696	52.0	10.0	7.8
8/1	6.0	132.0	6.5	15.5	6.5	70.0	61.0	14.0	11.0
8/8	11.5	56.0	8.5	14.5	2.0	35.5	40.0	12.0	5.0
8/15	38.5	74.0	14.0	26.0	8.5	9.0	41.5	14.5	14.8
8/22	18.0	13.0	8.5	19.5	4.5	481.5	20.0	11.5	4.8
8/29	36.5	13.5	11.0	36.5	1.5	107.5	12.0	11.0	8.0
9/6	33.5	38.5	18.5	35.5	1.0	73.0	13.5	10.0	7.3

[Section: GENERAL INFO]

SCAFFOLDS Fruit Journal

Index, Volume 25, 2016

No. 1, March 21

GENERAL INFO

New year of Scaffolds intro

NEWA weather station maintenance

DISEASES

Fungicide update for 2016

CHEM NEWS

Product registration update

No. 2, March 28

INSECTS

Early season oil strategies

San Jose Scale and White Prunicola Scale

No. 3, April 4

INSECTS

Dogwood borer and Black stem borer

DISEASES

Green tip and the NEWA Scab Model

No. 4, April 11

INSECTS

Pear Psylla

GENERAL INFO

New Degree Day tool in NEWA

CHEM NEWS

Erratum: Pear sections of Recommends

No. 5, April 18

INSECTS

Pre-bloom tarnished plant bug

Selecting/Using insecticide premixes

GENERAL INFO

Cornell Fruit Field Day 2016

No. 6, April 25

INSECTS

Pest prediction models on NEWA

DISEASES

Stone fruit disease management

GENERAL INFO

Cornell Fruit Field Day 2016

No. 7, May 2

INSECTS

Black stem borer emergence

Early Lepidoptera management

DISEASES

Rust diseases of apples and pears

GENERAL INFO

Cornell Fruit Field Day 2016

No. 8, May 9

CHEM NEWS

Nealta miticide NYS registration

Danitol 2(ee) for black stem borer

DISEASES

Fire blight management

GENERAL INFO

Cornell Fruit Field Day 2016

No. 9, May 16

INSECTS

Petal fall insects

DISEASES

Apple scab & quince rust in Hudson Valley

GENERAL INFO

Cornell Fruit Field Day 2016

No. 10, May 23

INSECTS

Orchard Radar Digest

Post-Petal fall insects

GENERAL INFO

Cornell Fruit Field Day 2016

No. 11, May 31

INSECTS

Orchard Radar Digest

Woolly apple aphid

Clearwing borers of stone fruits

GENERAL INFO

Cornell Fruit Field Day 2016

No. 12, June 6

INSECTS

Orchard Radar Digest

Codling moth management

San Jose scale and White Prunicola scale

GENERAL INFO

Cornell Fruit Field Day 2016

No. 13, June 13

INSECTS

Orchard Radar Digest

Potato Leafhopper

BMSB Guidance Document for Orchard Crops

GENERAL INFO

Cornell Fruit Field Day 2016

No. 14, June 21

INSECTS

Orchard Radar Digest

Summer foliar insects

Mite management

GENERAL INFO

Cornell Fruit Field Day 2016

No. 15, June 27

INSECTS

Orchard Radar Digest

Apple maggot

GENERAL INFO

Cornell Fruit Field Day 2016

No. 16, July 5

INSECTS

Orchard Radar Digest

Summer insects lineup

GENERAL INFO

Cornell Fruit Field Day 2016

No. 17, July 11

INSECTS

Orchard Radar Digest

Beneficial insects

GENERAL INFO

Cornell Fruit Field Day 2016

No. 18, July 18

INSECTS

Orchard Radar Digest

Comstock mealybug

CHEM NEWS

Magister SC miticide labeled in cherries

Bifenthrin Sect. 18 approved for BMSB

GENERAL INFO

Wayne Co. Fruitgrower Tour

No. 19, July 25

INSECTS

Orchard Radar Digest

Midsummer pest roundup

GENERAL INFO

Wayne Co. Fruitgrower Tour
Spanish-Speaking Fruit Summer Tour

No. 20, August 1

INSECTS

Orchard Radar Digest
August pest checklist
Dock sawfly

GENERAL INFO – EVENTS

Wayne Co. Fruitgrower Tour
Spanish-Speaking Fruit Summer Tour
IPM Climate & Weather Conference
Cornell Fruit Pest Control Field Day

No. 21, August 22

INSECTS

Orchard Radar Digest
Pest Focus

GENERAL INFO – EVENTS

Cornell Fruit Pest Control Field Days

No. 22, August 29

DISEASES

Preventing Bitter Rot In Apples

INSECTS

2016 Tree fruit arthropod pest review

Summary of 2016 Pest Events

GENERAL INFO – EVENTS

Cornell Fruit Pest Control Field Days

No. 23, September 6

DISEASES

Apple trunk problems can reduce profitability

INSECTS

2016 Insect trap catch summary

GENERAL INFO

Index of Scaffolds Volume 25, 2016

This material is based upon work supported by Smith Lever funds from the Cooperative State Research, Education, and Extension Service, U.S. Department of Agriculture. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.

Scaffolds is published weekly from March to September by Cornell University -- NYS Agricultural Experiment Station (Geneva), and Ithaca -- with the assistance of Cornell Cooperative Extension. New York field reports welcomed. Send submissions by 2 p.m. Monday to:

Scaffolds Fruit Journal

Editor: A. Agnello

Dept. of Entomology, NYSAES

630 W. North St.

Geneva, NY 14456-1371

Phone: 315-787-2341 FAX: 315-787-2326

E-mail: ama4@cornell.edu

Online at

<<http://www.scaffolds.entomology.cornell.edu/index.html>>